

ON HEREDITARY RADICALS OF SEMIGROUPS WITH ZERO

Ferenc Szász*

(Received October 7, 1974)

In this note, which can be considered as a continuation of author's paper [14], we shall verify that there exists a common generalization of some assertions of Robert Shulka [11], if we use general hereditary radicals of semigroups with zero, instead of various concrete radicals, e.g. Clifford's [2], Schwarz's [8] radicals, prime and completely prime radicals, respectively, used by R. Shulka. Our generalization also dualizes a result, asserted before for associative rings and alternative rings by author [13], on some join-endomorphisms of the lattice of all twosided ideals of the ring.

The fundamental notions for semigroups can be found in the book's A. H. Clifford and G. B. Preston [3] and E. S. Ljapin [5]. Various concrete radicals for semigroups were discussed e.g. by J. Bosak [1], A. H. Clifford [2], H. J. Hoehnke [4], J. Luh [6], S. Schwarz [8], H. Seidel [9], L. N. Shevrin [10] and author [12]. The possibility to investigate general radicals for semigroups with zero element was shown e.g. by author [14].

For a radical R and for an ideal I of the semigroup S , let $\phi(I)/I$ be the R -radical of S/I .

A radical R is called *hereditary*, if every two sided ideal I of an R -radical semigroup is R -radical.

Proposition 1. The mapping $I \rightarrow \phi(I)$, defined above for ideals of a semigroup, is monotone.

Proof. By D. Rees [7] the first and second isomorphism theorems hold for the factor semigroups. Assume $I_1 \subset I_2$ for the ideals $I_j (j=1, 2)$. Then we have:

$$(*) \quad \phi(I_1) \cup \phi(I_2) / \phi(I_2) \sim \phi(I_1) / \phi(I_1) \cap \phi(I_2).$$

Since here the right side is isomorphic to the factor semigroup $R(S/I_1) / (\phi(I_1) \cap \phi(I_2)) / I_1$ it is R -radical. At the same time, the left side of $(*)$ is a two sided ideal of $S/\phi(I_2) \sim (S/I_2) / (\phi(I_2)/I_2)$ hence it is, by Theorem 1 of author [14], also R -semisimple. This yields $\phi(I_1) / \phi(I_1) \cap \phi(I_2) = 0$

*Mathematical Institute of Hungarian Academy of Sciences, Budapest, Hungary.

consequently the desired inclusion $\phi(I_1) \subset \phi(I_2)$, indeed.

Proposition 2. For arbitrary radical R and two sided ideal I of the semigroup S , if I and S/I are R -radical, then also S is an R -radical semigroup.

Proof. Immediately follows, using the definition of the R -radical (see [14]) and that of Rees factor semigroups and isomorphism theorems (see [7]).

Theorem 3. For an arbitrary hereditary radical R of a semigroup S the mapping $I \rightarrow \phi(I)$ is a meet-endomorphism of the lattice of all two sided ideals of S .

Proof. By Proposition 1, we obviously have

$$\phi(I_1) \cap \phi(I_2) \supset \phi(I_1 \cap I_2).$$

Since

$$\phi(I_1) \cap \phi(I_2) / \phi(I_1) \cap \phi(I_2) \cap I_j \sim \phi(I_1) \cap \phi(I_2) \cup I_j / I_j \subset \phi(I_j) / I_j$$

and since R is hereditary, $\phi(I_1) \cap \phi(I_2) / \phi(I_1) \cap \phi(I_2) \cap I_j$ is R -radical, for $j=1$ and 2 .

Furthermore, e.g. also $\phi(I_1) \cap I_2 / I_1 \cap I_2$ is R -radical, since it can be isomorphically embedded as an ideal, by the first isomorphism theorem, into $\phi(I_1) / I_1$ and since R is hereditary. Similarly, it can be proved, that $I_1 \cap \phi(I_2) / I_1 \cap I_2$ is R -radical.

Now, since $I_j \cap \phi(I_k) / I_1 \cap I_2$ for $j \neq k$ and $j, k=1, 2$ and $\phi(I_1) \cap \phi(I_2) / I_j \cap \phi(I_k)$ are R -radical, by Proposition 2, also $\phi(I_1) \cap \phi(I_2) / I_1 \cap I_2$ is R -radical, which implies $\phi(I_1) \cap \phi(I_2) \subset \phi(I_1 \cap I_2)$ and hence $\phi(I_1) \cap \phi(I_2) = \phi(I_1 \cap I_2)$. Thus $I \rightarrow \phi(I)$ is a meet-endomorphism, indeed, which completes the proof.

References

- [1] J. Bosák, On radicals of semigroups (in Russian), *Matematicko-Fyzikalny Časopis SAV* 12, 12, 230-234 (1962).
- [2] A. H. Clifford, Semigroups without nilpotent ideals, *Amer. Jour.* 71, 46-58 (1949).
- [3] ——— and G. B. Preston, *The Algebraic Theory of Semigroups*, I., II. Providence (1961 and 1967).
- [4] H. J. Hoehnke, Structure of semigroups, *Canadian Jour. Math.* 18, 449-491 (1966).
- [5] E. S. Liapin, Semigroups (in Russian), Mosoow (1960).
- [6] J. Luh, On the concepts of radical of semigroup having kernel, *Portugaliae Math.* 19 189-198 (1960).
- [7] D. Rees, On semigroups, *Proc. Cambridge Philos. Soc.* 36, 387-400 (1940).
- [8] S. Schwarz, Über Theorie der Halbgruppen (in Slovakian), *Sbornik Prac Prirodovedecky Fakulty Slovenskij Univerzity v Bratislave* 6, 1-64 (1943).
- [9] H. Seidel, Über das Radikal einer Halbgruppe, *Math. Nachrichten* 29, 255-263 (1965).
- [10] L. N. Shevrin, On general theory of semigroups (in Russian), *Mat. Sbornik* 53 (95): 3, 367-586 (1961).
- [11] R. Shulka, On nilpotent elements, ideals and radicals of semigroups (in Russian),

Mat.
[12] F. S
cshe
(196
[13]
des
(196
[14]
(197

of the semigroup.

radical (see [7]).

roup S the sided ideals

- [12] Matematicko-Fyzikalny Časopis SAV 13: 3, 209-222 (1963).
F. Szász, Radikalbegriffe für Halbgruppen mit Nullelement, die dem Jacobson-
schen ringtheoretischen Radikal ähnlich sind, Math. Nachrichten 34, 157-161
(1967).
- [13] ———, Ein radikaltheoretischer Vereinigungsendomorphismus des Idealverban-
des der Ringe, Annales Univ. Budapest Roland Eotvos, Sectio Math. 12, 73-75
(1969).
- [14] ———, On radicals of semigroups with zero, I., Proc. Japan Acad. 46, 595-598
(1970).

ul, for $j=1$

can be iso-
morphism, into
that

$\phi(I_j \cap \phi(I_k))$
ical, which
(2). Thus
of.

ikalny Časopis

46-58 (1949).
roups, I., II.

49-491 (1966).

Portugaliae

(1940).

Prac Priro-

29, 255-263

Sbornik 53

(in Russian),